

Absolute properties of BG Ind – a bright F3 system just leaving the Main Sequence^{*}

M. Rożyczka,^{1†} J. Kaluzny,¹ W. Pych,¹ M. Konacki,²
K. Małek,^{3‡} L. Mankiewicz,^{3‡} M. Sokołowski,^{4‡} A.F. Żarnecki^{5‡}

¹*Nicolaus Copernicus Astronomical Center, Bartycza 18, 00-716 Warszawa, Poland*

²*Nicolaus Copernicus Astronomical Center, Rabiańska 8, 87-100 Toruń, Poland*

³*Center for Theoretical Physics PAS, Al. Lotników 32/46, 02-688 Warszawa, Poland*

⁴*The Andrzej Sołtan Institute for Nuclear Studies, Hoża 69, 00-681 Warszawa, Poland*

⁵*Institute for Experimental Physics, University of Warsaw, Hoża 69, 00-681 Warszawa, Poland*

Accepted ... Received ... in original form ...

ABSTRACT

We present photometric and spectroscopic analysis of the bright detached eclipsing binary BG Ind. The masses of the components are found to be 1.428 ± 0.008 and $1.293 \pm 0.008 M_{\odot}$ and the radii to be 2.290 ± 0.017 and $1.680 \pm 0.038 R_{\odot}$ for primary and secondary stars, respectively. Spectra- and isochrone-fitting coupled with colour indices calibration yield $[Fe/H] = -0.2 \pm 0.1$. At an age of 2.65 ± 0.20 Gyr BG Ind is well advanced in the main-sequence evolutionary phase - in fact, its primary is at TAMS or just beyond it. Together with three similar systems (BK Peg, BW Aqr and GX Gem) it offers an interesting opportunity to test the theoretical description of overshooting in the critical mass range $1.2 - 1.5 M_{\odot}$.

Key words: binaries: eclipsing – stars: fundamental parameters – stars: evolution

1 INTRODUCTION

The 6th magnitude star BG Ind was classified as an F3 dwarf by Malaroda (1975), and several years later established as an eclipsing binary by Manfroid & Mathys (1984) and Mathys, Manfroid & Renson (1986), who identified it as a detached system with partial eclipses and a period of 1.464047 d based on *ubvy* data from 1984. The first radial velocity measurements of BG Ind were performed by Andersen, Jensen & Nordström (1984). As they wrote, “The plates immediately showed spectral lines of two components of rather similar type, the lines of one component (the primary) being noticeably stronger and broader than those of the secondary.” Although their data were rather scarce (just two spectra), the resulting estimates of masses and radii of the components ($m_1 \sim 1.4 M_{\odot}$, $R_1 \sim 2.0 R_{\odot}$ and $m_2 \sim 1.2 M_{\odot}$, $R_2 \sim 1.5 R_{\odot}$) proved to be surprisingly accurate. Additional Strömgren photometry was collected in 1986 by Van Hamme & Manfroid (1988; hereafter VHM) who also solved the *vby* light curves of the system, and found an improved value of the period ($P = 1.464069$ d). For the fixed mass ratio $q = 0.85$ taken from Andersen et al. (1984)

they obtained $m_1 = 1.41 M_{\odot}$, $R_1 = 2.22 R_{\odot}$ and $m_2 = 1.20 M_{\odot}$, $R_2 = 1.60 R_{\odot}$.

Revised data of Mathys, Manfroid & Renson (1986) and VHM, together with additional points from 1987, were catalogued by Manfroid et al. (1991) and Sterken et al. (1993) as a part of the *Long Term Photometry of Variables* program conducted at ESO. BG Ind was monitored by Hipparcos satellite Perryman et al. (1997) and robotic telescopes ASAS (Pojmanski 2001) and Pi of the Sky (Małek et al. 2010); it is also included in *The Geneva-Copenhagen survey of the solar neighborhood* (Holmberg, Nordström & Andersen 2009). Recently, a spectroscopic solution of BG Ind based on data collected in 2006 has been published by Bakış et al. (2010), who for $P = 1.464069$ d and $i = 74.14^{\circ}$ found by VHM obtained $m_1 = 1.47 \pm 0.01 M_{\odot}$, $m_2 = 1.31 \pm 0.01 M_{\odot}$, and a semimajor axis $a = 7.64 \pm 0.04 R_{\odot}$.

With so much data available, BG Ind may seem to be well explored and hence of little interest. However, there are at least three reasons to investigate it more thoroughly than in the papers mentioned above. First, VHM encountered problems with phasing, and they recommended “further monitoring of this binary system for minimum times in order to obtain improved ephemeris, and to allow a study of the behavior of its orbital period”. Second, systemic radial velocity measurements gave diverging results: $v_{\gamma} = 39.8 \pm 4$

^{*} Based in part on data obtained at the South African Astronomical Observatory.

[†] E-mail: mnrc@camk.edu.pl

[‡] for the Pi of the Sky team

km s⁻¹ (VHM) and 59.4 ± 5 km s⁻¹ (Bakış et al. 2010). To make the confusion even larger, from three velocity components of BG Ind with respect to the Sun listed by Holmberg, Nordström & Andersen (2009) one obtains the *total* velocity of 20.3 km s⁻¹.

Third, and most important, both components of BG Ind belong to the interesting mass range of 1.1–1.5 $M_{\odot}$ in which convective cores begin to develop, affecting evolutionary tracks and isochrones via overshooting-related uncertainties. Moreover, their masses are remarkably similar to those of BK Peg – a main sequence binary recently studied by Clausen et al. (2010). Since the metallicity is similar in both cases, and smaller stellar radii observed in BK Peg ($R_1 = 1.987 \pm 0.008 R_{\odot}$, $R_2 = 1.473 \pm 0.017 R_{\odot}$) indicate a slightly less advanced evolutionary stage, these two systems are a potential source of valuable information concerning the endphases of hydrogen burning in stars somewhat more massive than the Sun. Bearing this in mind, we decided to re-analyze BG Ind with the aim to determine its precise physical parameters and evolutionary status.

Our paper is based on photometric and spectroscopic data described in Sect. 2. The analysis of the data is detailed in Sect. 3, and its results are discussed in Sect. 4.

2 OBSERVATIONAL MATERIAL AND DATA REDUCTION

2.1 Photometry

Our photometric solutions are based on six sets of photometric measurements listed in Table 1, spanning a period from May 1986 to November 2009 (points with $\Delta m > 0.1$ mag, where Δm stands for the deviation from the preliminary fit (see Sect. 3.2), were rejected from the original ASAS and Pi datasets). We found that all photometric data from Table 1 neatly phase with the ephemeris

$$\begin{aligned} t_0 &= \text{HJD } 47876.3792 \pm 0.0004 \\ P &= 1.46406335 \pm 0.00000002 \text{ d} \end{aligned} \quad (1)$$

(see Fig. A1), and we are rather confident that the period of BG Ind remained constant for over 23 years. The only indication suggesting a possible period change comes from the earliest observations of this system (runs 1 and 2 of VHM, not included in our analysis).

Available online are also Hipparcos B_T and V_T lightcurves (Perryman et al. 1997), and Strömgren u lightcurve of Manfroid et al. (1991) and Sterken et al. (1993). Their quality was too poor to use them for photometric solutions; however B_T and V_T data yielded a useful estimate of the temperature of the hotter component (see Sect. 3.2). We note here that according to (Suchkov, Makarov & Voges 2003) the Strömgren ($b - y$) excess of BG Ind amounts to 0.001 only, so that reddening effects can be neglected.

2.2 Spectroscopy

Because of problems with the systemic velocity of BG Ind mentioned in Sect. 1 we decided to use our own spectral data, collected in September / October 2007 with the fibered Giraffe spectrograph on the 1.9-m Radcliffe telescope at the South African Astronomical Observatory. The seeing

Table 1. List of light curves used in this paper for photometric solutions of BG Ind.

Label	Filter	1 st day [HJD-2400000]	last day	Number of data points	Ref.
MSb	Strömgren b	44582	47070	165	1,2
MSv	Strömgren v	46582	47070	172	1,2
MSy	Strömgren y	46582	47070	172	1,2
HipH	Hipparcos H	47888	49054	142	3
ASV	Johnson V	51997	55167	576	4
PiR	Cousins R	53902	54933	828	5

1: Manfroid et al. (1991); 2: Sterken et al. (1993); 3: Perryman et al. (1997); 4: Pojmanski (2001); 5: Małek et al. (2010). Pi of the Sky observations were done without filters, but the combined sensitivity of optical system and detector closely matched that of Cousins R-filter.

oscillated between 1.5 arcsec and 2.5 arcsec. A 2.7 arcsec entrance window provided a resolution of almost $R = 40000$ at $\lambda = 4470$ Å. During the observations pairs of scientific spectra were taken, separated by an exposure of a thorium-argon hollow-cathode lamp. The exposure times per spectrum ranged from 400 s to 900 s. The observations were reduced within the IRAF ¹ ECHELLE package. After bias and flat-field correction each pair of the frames was combined into a single frame, allowing for the rejection of cosmic ray hits. Altogether, 23 reduced spectra were obtained. For further analysis a wavelength range extending from 4300 Å to 5800 Å was used, in which most of the spectra had $20 < S/N < 40$ with a few cases of lower quality.

Our method of radial velocity measurements is based on the broadening function formalism introduced by Rucinski (2002) and it was described in detail by Kaluzny et al. (2006). ² The spectrum of the bright star HD 200163 (spectral type F3V), obtained in the same observing period and with the same instrumental setup, served as a template. The resulting barycentric radial velocities are listed in Table 2. The mean error, estimated from velocity-curve fitting, is ± 0.63 km s⁻¹.

The binary spectrum of BG Ind was disentangled using the code of Konacki et al. (2010) with the aim to estimate the temperature of the components. Unfortunately, because of strong rotational broadening and blending of the lines the only method we could apply was a direct comparison to synthetic spectra (see Sect. 3.2).

3 ANALYSIS

The components of BG Ind have masses in the range 1.1–1.5 $M_{\odot}$ (VHM, Bakış et al. 2010), so that their envelopes should be convective. Consequently, we adopted gravity darkening exponents $g_1 = g_2 = 0.32$ and bolometric albedos $A_1 = A_2 = 0.5$. Just in case, we also obtained a solution with radiative envelopes ($g_1 = g_2 = A_1 = A_2 = 1.0$) and

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the AURA, Inc., under cooperative agreement with the NSF.

² The software used in this paper is available at <http://users.camk.edu.pl/pych/BF/>.


Table 2. Barycentric radial velocities of BG Ind phased according to the ephemeris given by equation (1).

HJD-2454000	Phase	v_1 [km s $^{-1}$]	v_2 [km s $^{-1}$]
363.398889	0.833	134.234	-83.756
363.420815	0.848	127.712	-76.513
366.290933	0.808	142.702	-91.436
368.315269	0.191	-78.691	152.746
368.335724	0.205	-83.175	157.484
368.365740	0.225	-87.252	161.310
369.244426	0.825	136.888	-85.218
369.263211	0.838	131.761	-79.637
371.249086	0.195	-80.656	153.364
371.266822	0.207	-82.224	157.092
371.284471	0.219	-85.197	159.509
371.315563	0.240	-86.278	161.496
371.328299	0.249	-86.007	162.845
371.344954	0.260	-86.568	162.832
371.361949	0.272	-85.781	160.540
371.379009	0.283	-83.505	158.200
371.395454	0.295	-82.164	157.302
371.412749	0.306	-79.292	153.182
372.255302	0.882	112.450	-57.583
372.275909	0.896	104.275	-48.576
372.292410	0.907	96.605	-41.796
372.308893	0.918	89.945	-33.640
376.386726	0.221	144.054	-92.932

we found that even within unrealistically small formal error limits calculated by PHOEBE which are all “radiative” parameters but one did not differ from the “convective” ones. The exception was the radius of the primary R_1 which in the radiative case was by 5.6% smaller (we define primary as the *more massive* component).

Since BG Ind is a short-period system, and broadening-function fitting yielded $v \sin i$ values close to those expected from the synchronous rotation for the radii obtained by VHM, we assumed full synchronization of both components. The effects of reflection were included, and a logarithmic limb-darkening based on tables by Van Hamme (1993) was used as implemented in PHOEBE 031a.

According to Holmberg, Nordström & Andersen (2009) the system is slightly underabundant in metals ($[Fe/H] = -0.3$). However, their estimate is based on Strömgren photometry of the *total* light, and as such it cannot be 100% reliable. As for the temperatures of the components, VHM assumed $T_1 = 7000$ K for the primary, and obtained $T_2 = 6450$ K for the secondary from the fit. While these values are certainly reasonable, they may be too high for the radii obtained by VHM which indicate that at least the primary is about to leave the main sequence or even has left it. Moreover, upon comparing our phased velocity curve with phased light curves from Table 1 we found that the hotter component of BG Ind is the *secondary* (see Fig. 1).³ The same conclusion follows from the velocities measured by Bakış et al. (2010) which also nicely phase with the ephemeris (1), showing, however, a much larger scatter (the rms residual from

**Figure 1.** Velocity curve of BG Ind based on data contained in Table 2. Systemic velocity and rms residuum are equal to 31.2 ± 0.1 km s $^{-1}$ and 0.63 km s $^{-1}$ respectively. Phase 0 corresponds to the center of the *shallower* photometric minimum.

our fit to their data is 4.6 km s $^{-1}$, only slightly larger than 4.4 km s $^{-1}$ found in the original paper).

Thus, while searching for an observational estimate of the temperature of BG Ind components we decided to focus on the secondary. As it rotates significantly slower than the primary and must be less evolutionary advanced, we expected it to be a more or less normal main-sequence star to which available color-temperature calibrations could be reliably applied.

3.1 Preliminary solutions

We started the analysis from searching for solutions with T_2 equal to 7000, 6500 or 6000 K, and $[Fe/H]$ equal to 0.0 or -0.3. We used PHOEBE interface (Prša & Zwitter 2005) to the Wilson-Devinney code (Wilson & Devinney 1971), solving for all lightcurves simultaneously. The essential aim of preliminary calculations was to check how strongly in this parameter range vary the gravitational acceleration of the secondary g_2 and the contributions of the secondary to the total light in various wavebands at $\varphi = 0.25$. As detailed in the next Section, the first of those parameters was needed for an estimate of T_2 based on our spectra, while the remaining ones – for an independent estimate of T_2 based on calibrations of color indexes.

There is no indication for a nonzero eccentricity in either light or velocity curves, but just in case we included iterations of e . The solutions yielded $e = 0.0 \pm 0.0005$, and an almost constant $\log g_2 = 4.14 - 4.15$. The primary’s acceleration g_1 was also almost constant, but smaller by a factor of 2, i.e. appropriate for the beginning of the subgiant branch rather than for the main sequence.

3.2 The temperature of the secondary

With g_2 fixed, we generated an array of synthetic spectra for $6000\text{K} \leq T \leq 7000\text{K}$ and $-1 \leq [Fe/H] \leq 0$ based on the library of (Coelho et al. 2005). The respective spacings ΔT and $\Delta[Fe/H]$ were equal to 100 K and 0.1. All spectra were rotationally broadened with $v \sin i = 78$ km s $^{-1}$ for the primary and 53 km s $^{-1}$ for the secondary, consistently with the mean values of i and component radii obtained from preliminary fits (73° , $2.35 R_\odot$ and $1.61 R_\odot$, respectively). Next,

³ The misidentification of VHM is entirely understandable when one remembers that they had just two velocity measurements and only an approximate value of the period.

Table 3. PHOEBE solutions with $T_2 = 6650$ K and $[Fe/H] = -0.2$.

light curve	rms [mmag]	i [deg]	a [$R_\odot$]	m_1 [$M_\odot$]	m_2 [$M_\odot$]	R_1 [$R_\odot$]	R_2 [$R_\odot$]	T_1 [K]	L_1 [$L_\odot$]	L_2 [$L_\odot$]
MSb	7.0	72.95	7.567	1.428	1.293	2.313	1.651	6312	7.37	4.62
MSv	7.5	72.95	7.567	1.428	1.293	2.303	1.663	6320	7.34	4.69
MSy	7.9	73.00	7.565	1.427	1.292	2.165	1.796	6443	7.00	5.47
HipH	6.4	73.12	7.560	1.424	1.289	2.226	1.723	6382	7.13	5.04
ASV	16.1	72.72	7.577	1.434	1.298	2.395	1.637	6375	8.22	4.55
PiR	11.4	73.03	7.567	1.428	1.293	2.340	1.610	6262	7.30	4.39

for each synthetic spectrum the sum of squared deviations from its observed counterpart was calculated. In the case of the secondary the smallest sum was obtained for $T_2 = 6500$ K and $[Fe/H] = -0.2$ (see Fig. A2 for the comparison of best-fitting spectrum to the observed one). In the case of the primary no unique minimum was found, for what broad and/or blended lines are probably to be blamed (e.g. MgI lines at 5167Å and 5173Å were observed as a single spectral feature). We estimate the uncertainties of temperature and metallicity at ± 100 K and ± 0.1 , respectively.

In all preliminary solutions with $[Fe/H] = 0$ the secondary to primary luminosity ratio L_2/L_1 (found from the contribution of each component to the total light of the system at $\varphi = 0.25$) was almost constant in each of b , v and y bands, amounting respectively to 0.570-0.573, 0.598-0.606 and 0.551-0.553. Given the total apparent magnitudes of BG Ind at $\varphi = 0.25$, we used these ratios to calculate the corresponding apparent magnitudes of the secondary, and we found its temperature from $(b - y) - T_{eff}$ calibration of Casagrande et al. (2010). The result for $[Fe/H] = 0.0$ was a remarkably constant $T_2 = 6523 - 6528$ K. Solutions with $[Fe/H] = -0.3$ were only slightly more diverging, with $T_2 = 6477 - 6487$ K. After averaging, we got $T_2 = 6525 \pm 63$ K and $T_2 = 6483 \pm 63$ K, respectively, where the error includes inaccuracies of color-temperature calibration and magnitude measurement at $\varphi = 0.25$. Based on the calibrations of Holmberg, Nordström & Andersen (2007), we also derived the observed $[Fe/H]$ index, which varied between -0.4 and -0.7. While being only marginally consistent with the lower of the $[Fe/H]$ values assumed in preliminary fitting, it confirmed that BG Ind is indeed metal-deficient.

The third estimate of T_2 was based on the Hipparcos photometry. Proceeding as before, we found the light ratio L_2/L_1 at $\varphi = 0.25$ in B_T and V_T bands, and, using $(B - V)_T - T_{eff}$ calibration of Casagrande et al. (2010), we derived $T_2 = 6650 \pm 83$ K for $[Fe/H] = 0$ and $T_2 = 6496 \pm 83$ K for $[Fe/H] = -0.3$.

For the fourth temperature estimate we used Hipparcos parallax of BG Ind ($\pi = 15.04 \pm 1.1$ mas) and ASAS photometry in V -band. Proceeding as before and using bolometric corrections of Lejeune, Cuisinier & Buse (1998), we found the observed V -band magnitude of the secondary ($V_2^o = 7.22$), and compared it to apparent magnitudes V_2^c calculated from preliminary solutions. Solutions with $T_2 = 6000$ which yielded $7.59 \leq V_2^c \leq 7.88$ were clearly ruled out. Those with $T_2 = 6500$ K and $T_2 = 7000$ K produced, respectively, $7.27 \leq V_2^c \leq 7.56$ and $6.95 \leq V_2^c \leq 7.24$, i.e. magnitudes almost consistent with V_2^o , but remaining on the low side in the first case and on the high side in the second

case. A nearly ideal agreement between observed and calculated magnitude was reached in an additional preliminary fit with $T_2 = 6750$ K, for which the difference $V_2^o - V_2^c$ was equal to just 0.02 mag.

Thus, the revised Hipparcos parallax pulled the temperature estimate firmly upward. Trusting it, we adopted $T_2 = 6650$ K with a fiducial uncertainty range of ± 100 K, marginally compatible with all available data. As it was detailed above, the trend observed in preliminary solutions indicated that the metallicities obtained from Strömgren photometry would cause T_2 to fall too low to be compatible with V_2^o . Bearing this in mind, we set $[Fe/H]$ to -0.2, i.e. to the value resulting from the spectroscopic estimate.

3.3 Final model fitting

For the final analysis the JKTEBOP code (Etzel 1981; Popper & Etzel 1981; Southworth, Maxted & Smalley 2004; Southworth et al. 2004) and a spectroscopic data solver written and kindly provided by G. Torres were used except PHOEBE. While JKTEBOP gives realistic estimates of the errors, it cannot be used when the stars are too distorted. Unfortunately, this is the case of BG Ind, where the primary's distortion exceeds the allowable limit by 10%. Consequently, we employed JKTEBOP solely to estimate the errors of the photometric solution (note that differences between the errors of the slightly inaccurate JKTEBOP solution and the errors of the appropriate solution must be small quantities of the second order which can be neglected). Similarly, realistic errors of the spectroscopic solution were found using the Torres code. (Because the Torres code requires center of mass velocities on input, it was necessary to correct the observed light center velocities for effects caused by the distortion of the components. Additive phase-dependent corrections were calculated using the Wilson-Devinney code; their values ranged from -0.86 to 0.27 km s⁻¹).

JKTEBOP cannot deal with multiple lightcurves, so that the final model fitting had to be performed separately for each of the photometric datasets listed in Table 1. We fixed $e = 0$ as indicated by preliminary solutions, and started the final calculations from PHOEBE fits which produced six sets of parameters listed in Table 3 whose second column shows the rms deviation of observed points from the fit. Next, we calculated errors of inclination i and relative radii of the components $r_{1,2} = R_{1,2}/A$ using JKTEBOP, and errors of $m_{1,2} \sin^3 i$ and $A \sin i$ using the Torres code. The errors were then transformed into errors of parameters returned by PHOEBE, and assigned to respective PHOEBE solutions. Finally, weighted averages of PHOEBE param-




Figure 2. Location of BG Ind components on $M - R$, $M - \log L$ and $\log T_{eff} - \log L$ planes. The lines are Dartmouth $[Fe/H] = -0.2$ isochrones for 2.60 and 2.67 Gyr (i.e. for extreme ages still fitting the primary on the $M - R$ plane).

ters and the errors of those averages were found from the standard formulae

$$\langle x \rangle = \frac{\sum_{j=1}^6 x_j / \sigma_j^2}{\sum_{j=1}^6 1 / \sigma_j^2}, \quad (2)$$

and

$$\sigma^2 = \frac{1}{\sum_{j=1}^6 1 / \sigma_j^2}. \quad (3)$$

The final parameters with their errors are listed in Table 4.

4 DISCUSSION AND CONCLUSIONS

In Sect. 1 we outlined three problems which prompted us to analyze BG Ind: variability of the period, doubtful systemic velocity and lack of accurate parameters of the system. As mentioned in Sect. 2.1, we found that between 1986 and 2009 the period remained constant at $1.46406335 \pm 0.00000002$ d. The only indication that it might have changed comes from the earliest observations of BG IND from 1981 and 1984 (runs 1 and 2 of VHM) which were not included in our data. As for the systemic velocity, our value of 31.2 ± 0.2 km s⁻¹ does not agree with any of those obtained by other authors. The value found by VHM (39.8 ± 4 km s⁻¹) was based on two measurements only, and its error was likely underestimated. 59.4 ± 5 km s⁻¹ of Bakış et al. (2010) is rather large for an F-type star in Sun's vicinity. The origin of such a large discrepancy with all remaining estimates is difficult to explain - we may only note that they did not observe any radial velocity standards, and it is conceivable that they reversed the sign of the heliocentric correction while reducing the data. Finally, the low $v_{total} = 20.3$ km s⁻¹ of Holmberg, Nordström & Andersen (2009) can be explained by the fact that they did not detect the binarity of BG Ind (this system is not included in their Table 2), and their velocity must have been contaminated by the orbital motion.

The parameters found in Sect. 3.3 are accurate enough for isochrone fitting. We used solar-scaled Dartmouth

Table 4. Physical parameters of BG Ind.

Parameter	Value			Unit
P	1.46406335	$\pm$	0.00000002	[d]
i	72.96	$\pm$	0.23	[deg]
e	0.000	$\pm$	0.0005	
a	7.567	$\pm$	0.013	$R_{\odot}$
m_1	1.428	$\pm$	0.008	$M_{\odot}$
m_2	1.293	$\pm$	0.008	$M_{\odot}$
R_1	2.290	$\pm$	0.017	$R_{\odot}$
R_2	1.680	$\pm$	0.038	$R_{\odot}$
$\log T_1$ [K]	3.803	$\pm$	0.018	
$\log T_2$ [K]	3.823	$\pm$	0.015	
$\log L_1 / L_{\odot}$	0.87	$\pm$	0.07	
$\log L_2 / L_{\odot}$	0.68	$\pm$	0.08	
$[Fe/H]$	-0.2	$\pm$	0.1	
age	2.65	$\pm$	0.20	Gyr

isochrones (Dotter et al. 2008) which include core overshooting defined as a product of the pressure scale height and a factor α_{over} which depends on stellar mass and composition (at nearly-solar metallicities it grows from 0.05 for $1.2 \leq M \leq 1.3 M_{\odot}$ through 0.1 for $1.3 < M \leq 1.4 M_{\odot}$ to 0.2 for $M > 1.4 M_{\odot}$). The convection itself is treated according to the standard mixing length theory with solar-calibrated mixing length parameter $\alpha_{ml} = 1.938$.

What makes BG Ind particularly interesting is that the masses of the components fall at the beginning and at the end of the range where α_{over} ramps up. First, we fitted isochrones calculated for $-0.4 \leq [Fe/H] \leq 0$ with a step of 0.1 to the most accurately determined parameters, i.e. masses and radii of the components. The respective ages we found were 2.27-2.43, 2.45-2.55, 2.60-2.67, 2.77-2.85, and 2.96-3.10 Gyr. Next, we checked how well these isochrones perform on $M - \log L$ and $\log T_{eff} - \log L$ planes. The best agreement was obtained for $[Fe/H] = -0.2$. The fit with $[Fe/H] = -0.1$ was almost equally good; that with $[Fe/H] = -0.3$ was still acceptable; and those with $[Fe/H] = 0$ and $[Fe/H] = -0.4$ had to be rejected. Thus, the isochrone fitting confirmed our spectroscopic estimate of




Figure 3. Systems with similar evolutionary advancement as BG Ind. The lines are Dartmouth $[Fe/H] = 0.0$ isochrones for 2.60 and 2.80 Gyr (BK Peg); 2.10 and 2.45 Gyr (BW Aqr); 2.45 and 2.70 Gyr (GX Gem).

metallicity which we finally fixed at $[Fe/H] = -0.2 \pm 0.1$. The corresponding age is 2.65 ± 0.20 Gyr.

The location of BG Ind on $M - R$, $M - \log L$ and $\log T_{eff} - \log L$ planes is shown in Fig. 2 together with $t = 2.60$ Gyr and $t = 2.67$ isochrones obtained for $[Fe/H] = -0.2$. One can see that the more massive primary has almost reached the beginning of the subgiant branch, while the secondary is still on its way to TAMS. The agreement between theoretical and observational data would be ideal if it were not for small discrepancies in R_2 and T_1 (by $\sim 3\%$ and $\sim 1.5\%$, respectively). The first one could originate from the fact that the eclipses of BG Ind are partial and the secondary is by almost 40% smaller than the primary (and therefore less deformed). As a result, and because of rather

poor quality of available photometric data, the accuracy of R_2 determination has to be markedly lower than that of R_1 . The second discrepancy, at a first glance rather insignificant, turned our attention because it occurs precisely where the effects of overshoot-treatment should be largest (the primary of BG Ind is a star with $M > 1.4M_\odot$ at TAMS). We decided to check if the same effect appears in other systems with similar masses and in similar evolutionary phase.

Based on a recent compilation of Clausen et al. (2010), we chose BW Aqr, BK Peg, and GX Gem whose component masses range from 1.26 to $1.49 M_\odot$, and whose $[Fe/H]$ indexes are consistent with 0. The fitting of solar-scaled Dartmouth isochrones for $[Fe/H] = 0$ yielded respective ages of 2.15 - 2.45, 2.6 - 2.8 and 2.50 - 2.75 Gyr, placed roughly

halfway between Yonsei-Yale and VRSS ages quoted in Table 12 of Clausen et al. (2010). Figs. 2 and 3 demonstrate that the temperature discrepancy, absent in the relatively unevolved secondary components of BG Ind and BK Peg, increases with the evolutionary advancement (whose best indicator is the distance from the sharp upturn of the isochrones on the $M - \log L$ plane) until it becomes clearly visible in GX Gem whose both components are at TAMS or have already left the main sequence. Note that luminosity errors are larger than those quoted by Clausen et al. (2010) - this is because we recalculated them according to the formula

$$\delta \log L = \sqrt{\left(2 \frac{\delta R}{R}\right)^2 + \left(4 \frac{\delta T}{T}\right)^2} \quad (4)$$

to make them consistent with ours.

The discrepancy is a 1- σ effect and as such it may not be real, however the trend it exhibits suggests there might be some physics behind. It is beyond the scope of our paper to identify physical factors or assumptions potentially responsible for such effect. Whether anybody decides to look for them or not, improving the quality of photometric and spectroscopic solutions of all four systems is certainly a worthwhile task, although in the case of BG Ind it might prove rather difficult because of strong rotational broadening.

ACKNOWLEDGMENTS

We thank Guillermo Torres for providing the spectroscopic data solver. Research of JK is supported by Foundation for Polish Science through the grant MISTRZ, and Polish Ministry of Science and Higher Education (PMSHE) through the grant N N203 379936 which also supports WP and MR. MK acknowledges the support from Foundation for Polish Science through the FOCUS grant and fellowship, and European Research Council through the Starting Independent Researcher Grant. Pi of the Sky is financed by PMSHE in 2009-2011 as a research project.

REFERENCES

- Andersen J., Jensen K.S., Nordström B., 1984, IBVS, 2642
 Bakış V., Bakış H., Bilir S., Soyduğan F., Soyduğan E., Demircan O., Eker Z., Yaz E., Tüysüz M., Şenyüz T., 2010, New Astr., 15, 1
 Casagrande L., Ramírez I., Meléndez J., Bessell M., Asplund M., 2010, A&A, 512, A54
 Clausen J.V., Frandsen S., Bruntt H., Olsen E.H., Helt B.E., Gregersen K., Juncher D., Krogstrup P., 2010, A&A, 516, 42
 Coelho P., Barbuy B., J. Meléndez J., Schiavon R.P., Castilho B.V., 2005, A&A, 443, 735
 Dotter A., Chaboyer B., Jevremović D., Kostov V., Baron E., Ferguson J.W. 2008, ApJS, 178, 89
 Etzel, P.B. 1981, NATO ASI Series C., Vol. 69, 111
 Holmberg J., Nordström B., Andersen J., 2007, A&A, 475, 519
 Holmberg J., Nordström B., Andersen J., 2009, A&A, 501, 941
 Kaluzny J., Pych W., Rucinski S.M., Thompson I.B., 2006, AcA, 56, 237
 Konacki M., Muterspaugh M.W., Kulkarni S.R., Helminiak K.G., 2010, ApJ, 719, 1293
 Lejeune T., Cuisinier F., Buser R.C., 1998, A&A Suppl. 130,65
 Malaroda S., 1975, AJ, 80, 637
 Małek K., Batsch T., Czyrkowski H., Ćwiok M., Dąbrowski R., Dominik W., Kasprowicz G., Majcher A., Majczyna A., Mankiewicz L., Nawrocki K., Pietrzak R., Piotrowski L.W., Ptasńska M., Siudek M., Sokoowski M., Użycki J., Wawer P., Wawrzaszek R., Wrochna G., Zaremba M., Aleksander F. Żarnecki A.F., 2010, Adv.Ast., 2010, 194946
 Manfroid J., Mathys G., 1984, IBVS, 2616
 Mathys G., Manfroid J., Renson P., 1986, A&A Suppl. 63, 403
 Manfroid J., Sterken C., Bruch A., Burger M., de Groot M., Duerbeck H.W., Duemmler R., Figer A., Hageman T., Hensberge H., Jorissen A., Madejsky R., Mandel H., Ott H.-A., Reitermann A., Schulte-Ladbeck R.E., Stahl O., Steenman H., Vander Linden D., Zickgraf F.-J., 1991, A&A Suppl., 87, 481
 Nordström B., Mayor M., Andersen J., Holmberg J., Pont F., Jørgensen B.R., Olsen E.H., Udry S., Mowlavi N., 2004, A&A, 418, 989
 Perryman M.A.C., Lindegren L., Kovalevsky J., Hoeg E., Bastian U., Bernacca P.L., Crézé M., Donati F., Grenon M., van Leeuwen F., van der Marel H., Mignard F., Murray C.A., Le Poole R.S., Schrijver H., Turon C., Arenou F., Froeschlé M., Petersen C.S. 1997, A&A, 323, L49
 Pojmanski G., 2001, ASPC, 246, 53
 Popper D.M., Etzel, P.B., 1981, AJ, 86, 102
 Prša A., Zwitter T., 2005, AJ, 628, 426
 Rucinski S.M., 2002, AJ, 124, 1746
 Suchkov A.A., Makarov V.V., Voges W., 2003, ApJ, 595, 1206
 Southworth J., Maxted P.F.L., Smalley B., 2004, MNRAS, 351, 1277
 Southworth J., Zucker S., Maxted P.F.L., Smalley B., 2004, MNRAS, 355, 986
 Sterken C., Manfroid J., Barzewski A.K., Bibó E., Bruch A., Burger M., Duerbeck H.W., Duemmler R., Heck A., Hensberge H., Hiesgen M., Inklaar F., Jorissen A., Juetner A., Kinkel U., Liu Zongli, Mekken M.V., Ng Y.K., Niarchos P., Puttmann M., Szeifert T., Spiller F., van Dijk R., Vogt N., Wanders I., 1993, A&A Suppl., 102, 79
 Van Hamme W., 1993, AJ, 106, 2096
 Van Hamme W., Manfroid J., 1988, A&A Suppl., 74, 247
 Wilson R.E., Devinney, E.J., 1971, AJ, 166, 605

APPENDIX




Figure A1. BG Ind light curves used in this paper (see Table 1 for the list), phased with the ephemeris (1). Individual curves are normalized to magnitude 0 at maximum light.




Figure A2. A section of the disentangled spectrum of the secondary (line) compared with the best fitting synthetic spectrum, obtained for $T_{eff} = 6500$ K, $g = 4.15$, $[Fe/H] = -0.2$, and rotationally broadened with $v \sin i = 53$ km s $^{-1}$ (dots). The largest differences appear in deep lines and in overlap regions of echelle orders.